

ORIGINAL ARTICLE

Untreated and enzyme-modified bovine whey products reduce association of *Salmonella* Typhimurium, *Escherichia coli* O157:H7 and *Cronobacter malonaticus* (formerly *Enterobacter sakazakii*) to CaCo-2 cells

R.M. Halpin¹, D.B. Brady¹, E.D. O’Riordan² and M. O’Sullivan²

¹ School of Biomolecular and Biomedical Sciences, University College Dublin, Belfield, Dublin, Ireland

² School of Agriculture, Food Science and Veterinary Medicine, University College Dublin, Dublin, Ireland

Keywords

association, CaCo-2 cells, *C. malonaticus*, *E. coli* O157:H7, *Salm.* Typhimurium, whey.

Correspondence

Rachel Halpin, School of Biomolecular & Biomedical Science, Ardmore House, University College Dublin, Dublin 4, Ireland.
E-mail: rachel.halpin@ucd.ie

2009/0179: received 29 January 2009,
revised 29 April 2009 and accepted 28 May
2009

doi:10.1111/j.1365-2672.2009.04436.x

Abstract

Aims: Adhesion of a micro-organism to a cell surface is often considered to be the first step in pathogenesis. Inhibiting this process may have therapeutic effects *in vivo*. This study investigates the inhibitory effects of various bovine whey products on the association of *Salm.* Typhimurium, *E. coli* O157:H7 and *C. malonaticus* (formerly *Enterobacter sakazakii*) to the human CaCo-2 cell line. Invasion of CaCo-2 cells by *Salm.* Typhimurium and *C. malonaticus* was also examined.

Methods and Results: Infection assays were performed by incubating pathogenic bacteria with CaCo-2 cells in the presence of untreated (UT) or enzyme-modified (EM) whey products. Associated micro-organisms were directly quantified by plate counts. Invasion of CaCo-2 cells by *Salm.* Typhimurium and *C. malonaticus* in the presence/absence of test materials was also quantified using gentamicin protection assays. At a concentration of 40 mg ml⁻¹, some UT whey products reduced association and invasion, but this effect was enhanced following hydrolysis with porcine pancreatic lipase.

Conclusions: Both UT and EM sweet whey protein concentrates (WPCs) were found to be particularly effective inhibitors of association and invasion. All EM whey products significantly ($P < 0.05$) inhibited invasion of *C. malonaticus* into epithelial cells, causing a 2-log reduction in the quantity of these micro-organisms internalized.

Significance and Impact of the Study: The present study suggests that whey products can inhibit association to and invasion of CaCo-2 cells by selected micro-organisms and may be useful in the treatment and/or prevention of foodborne infections.

Introduction

Food consumed by humans is rarely sterile. Micro-organisms present in food can lead to spoilage and/or foodborne illness, with the latter causing millions of cases of infection and in some cases even death every year (Meng and Doyle 1998). *Salmonella enterica* serovar Typhimurium (*Salm.* Typhimurium) and *Escherichia coli* O157:H7 (enterohaemorrhagic *E. coli*, EHEC) have been recognized as pathogens for many years, but it is only in the last

30 years that these bacteria have been considered to be predominantly foodborne (Meng and Doyle 1998). Salmonellosis is a zoonotic infection, with infected animals being a major source of illness (Bezirtzoglou *et al.* 2000), and infection in humans is often because of the consumption of undercooked poultry, eggs or egg-containing foods (Rodrigue *et al.* 1990). *Salm.* Typhimurium is considered to be one of the most common causes of salmonellosis worldwide (WHO, 2005). *Escherichia coli* is an inhabitant of the gut (both humans and animals),

but some strains are pathogenic (Bezirtzoglou *et al.* 2000). *Escherichia coli* O157:H7 is a clinically important food pathogen that can cause haemorrhagic colitis and haemolytic uremic syndrome (HUS) (Gu *et al.* 2008). EHEC has an extremely low infectious dose, and it is estimated that as few as 100 cells is adequate to cause infection (Kaper *et al.* 2004). *Enterobacter sakazakii* was listed as a new species in 1980, but a taxonomic reclassification of this micro-organism has been proposed, as *E. sakazakii* has been found to consist of five species within a new genus now referred to as 'Cronobacter' (Iversen *et al.* 2007). *Cronobacter malonaticus* is one such species present within this new genus. This bacterium is described as an emerging opportunistic pathogen, which can cause local necrotizing enterocolitis, systemic bacteremia and meningitis (Kim and Loessner 2008). *Cronobacter* spp. has been isolated from milk powder, cheese, sausage meat, vegetables, bread, herbs and spices (Mullane *et al.* 2007; Gurtler *et al.* 2005 and Kandhai *et al.* 2004), but powdered infant formula (PIF) is considered to be a major vehicle of transmission, with neonates being most at risk of infection (Kim and Loessner 2008).

Once ingested, micro-organisms present in contaminated foods can adhere to the host's cell surfaces. Adhesion of a microbe to a cell surface is considered to be the first step of pathogenesis (Finlay and Falkow 1997). Lectins on the surface of bacteria adhere to specific receptors on epithelial cells of the intestinal tract (Nakajima *et al.* 2005). Some pathogenic micro-organisms are capable of entering and surviving within epithelial cells following initial adherence, a process known as invasion (Finlay and Falkow 1997). Cultured eukaryotic cell lines are a common method employed in the study of bacterial adherence and invasion as they provide researchers with reproducible and less complicated infection models (Tang *et al.* 1993). One of the most extensively used is the CaCo-2 cell line, which was isolated from human colon carcinoma (Fogh *et al.* 1977). These cells, under standard culture conditions, differentiate to produce monolayers expressing characteristics of mature enterocytes (Pinto *et al.* 1983).

Blocking the initial adherence of foodborne pathogens to intestinal epithelial cells may be a suitable approach to preventing occurrence of infections (Nakajima *et al.* 2005). Food components capable of inhibiting initial adherence are promising agents of intervention. Several carbohydrate components of food have exhibited a positive effect against intestinal infection (Nakajima *et al.* 2005), such as sialylated oligosaccharides from bovine or human milks (Sugita-Konishi *et al.* 2002). Recently, whey and dairy products have been shown to reduce the adherence of the dental caries causing bacterium *Streptococcus mutans* to hydroxylapatite, an analogue of tooth enamel

(Halpin *et al.* 2008). Whey was once considered to be a waste product of the cheese-making process, but in recent years has had its status upgraded to coproduct and is described as a 'functional food' (Marshall 2004). Many studies have shown enzymatic hydrolysis of whey proteins produces a plethora of peptides, exhibiting a wide array of bioactive properties (Meisel 1998).

The objective of this study was to examine the influence of a variety of whey products on the interaction of *Salm. Typhimurium*, *E. coli* O157:H7 and *Cronobacter malonaticus* with CaCo-2 cells. The effect of pretreating the whey products with PPL (porcine pancreatic lipase) on any such influence was also examined.

Materials and methods

Source and analysis of dairy powders

Sweet whey protein concentrate (WPC, 80% protein), acid WPC 80 (AWPC80), sweet WPC 35 (SWPC35), whey protein isolate (WPI), whey powder (WP) and demineralized whey (DW) powders were supplied by Carbery Milk Products (Ballineen, Cork, Ireland). Albumin from chicken egg white (grade V) was supplied by Sigma (Poole, Dorset, UK).

Compositional analysis was performed on each whey product using standard methods. Ash content was determined according to the methods of Malkomesius and Nehring (1951). Fat content was determined according to the method of Röse-Gottlieb (International Dairy Federation (IDF) 1987), protein content was determined by the Kjeldahl method (IDF 1993a), and the moisture content was determined by oven drying (IDF 1993b).

Hydrolysis conditions

Crude porcine pancreatic lipase (PPL, Sigma) containing 100–400 units mg⁻¹ protein was used throughout the study. Hydrolysates were prepared in a Fercmac 200 fermentor (Electrolab Ltd, Tewkesbury, UK) as follows: c. 2% (w/v) solution of substrate was prepared by dissolving 20 g of whey product in 900 ml of sterile-distilled water and heating to 37°C with stirring for 30 min. Lipase solution (1 g of PPL in 100 ml of sterile H₂O) was added to the substrate solution to give a final incubation volume of 1 l. The hydrolysates were then incubated for 2 h at 37°C with stirring. Following this, hydrolysates were heated at 60°C for 10 min in order to denature the enzyme(s). Each hydrolysate was placed on ice and allowed to cool to below 10°C (c. 45 min), before being frozen using liquid nitrogen and subsequently lyophilized (Moduloyo, Edwards High Vacuum, Manor Royal, Crawley, Sussex, UK).

Bacteria and growth conditions

Salmonella Typhimurium (ATCC 14028) and *E. coli* O157:H7 (ATCC 43888) were obtained from the American Type Culture Collection (Rockville, MD, USA), and *C. malonaticus* (DSM 18702) was obtained from the German Collection of Micro-organisms and Cell Cultures (DSMZ, Braunschweig, Germany). Primary cultures were grown overnight in 10 ml of Luria-Bertani (LB) broth (Sigma) at 37°C. Then, 1% inoculum was prepared by adding 400 µl of overnight culture to 39.6 ml of fresh prewarmed LB broth before re-incubating at 37°C. *Salm.* Typhimurium, and *E. coli* O157:H7 were grown to mid-log phase. *Cronobacter malonaticus* was grown to late-log phase, as it has been recently reported that adherence of *E. sakazakii* is at its maximum at this stage of growth, after at least 4 h of culturing (Mange *et al.* 2006). Bacteria were then collected by centrifugation at 3220 g (Eppendorf 5810R, Cambridge, UK) for 10 min and were resuspended in supplement-free Dupleco's Modified Eagle's Medium (DMEM, Gibco), so that the final concentration of bacteria could be *c.* 10⁸ cells ml⁻¹.

CaCo-2 cell culture

CaCo-2 cells were obtained from the European Collection of Cell Cultures (ECACC, Wiltshire, UK). At late confluency, these cells express both structural and functional characteristics of enterocytes present in the small intestine (Hendricks *et al.* 1996). Cells were routinely cultured in DMEM supplemented with 10% heat-inactivated foetal bovine serum (FBS), 1% nonessential amino acids 100X, 1% penicillin/streptomycin solution and 1% fungizone containing 250 µg ml⁻¹ amphotericin B. All supplements were supplied by Gibco. The cells were initially grown in T75cm² flasks (Sarstedt, Nümbrecht, Germany) and upon confluency (*c.* 1.5 × 10⁶ cells ml⁻¹, 7–10 days) were passaged using 0.25% trypsin (Gibco). For infection assays, monolayers were cultured in 24-well tissue culture plates (Sarstedt). CaCo-2 cells were seeded at a density of 5 × 10⁵ cells well⁻¹, and growth medium was changed every other day. These cells are known to be fully differentiated after being cultured for 19 days (Koninkx 1995). Maintenance of cells and subsequent experiments were carried out at 37°C in 5% CO₂/95% air atmosphere (Binder Apt Line C150, Tuttlingen, Germany), between passage numbers 37 and 55, with *c.* 10⁶ CaCo-2 cells well⁻¹.

Viability of bacterial and epithelial cells

The trypan blue (Sigma) dye exclusion test was used to determine whether test materials affected the viability of CaCo-2 cells. Test materials were prepared in

supplement-free DMEM (SFM) and added to wells containing monolayers, followed by incubation for 1 h at 37°C and 5% CO₂. The monolayers were then washed twice with SFM, before being trypsinized and added to 0.4% trypan blue (1 : 1). An inverted light microscope (Ceti, Kontich-Antwerp, Belgium) was used to examine cells to determine viability. Viable epithelial cells exclude trypan blue while dead cells allow entry and appear blue when viewed under the microscope. The percentage viability was calculated as follows:

$$100 - (\text{Number of dead cells} / \text{Total Number of cells} \times 100)$$

Viability of bacteria was determined by direct contact studies, where each micro-organism was incubated with test materials under typical assay conditions and subsequently enumerated by spread plates following appropriate dilution. As whey products are not sterile, selective agars were used to quantify the number of pathogenic bacteria (i.e. to eliminate any 'background count' because of micro-organisms such as lactobacilli). Brilliant green, McConkey and Chromogenic *Enterobacter sakazakii* agars (DFI formulation) (Oxoid, Hampshire, UK) were used to selectively cultivate *Salm.* Typhimurium, *E. coli* O157:H7 and *Cronobacter malonaticus*, respectively.

Infection assays

Prior to infection assays, CaCo-2 cells were washed twice in sterile phosphate-buffered saline (PBS, Sigma) to remove traces of antibiotic and equilibrated in SFM at 37°C and 5% CO₂ for at least 2 h. (i) Association and (ii) invasion of pathogenic bacteria were examined as follows:

Quantification of association

Aliquots (500 µl) of test material dispersed in SFM (80 mg ml⁻¹) or 500 µl of SFM (to act as a negative control) were added to each well, followed by 500 µl of pathogens suspended in SFM (*c.* 10⁸ CFU ml⁻¹). The final concentration of test material was 40 mg ml⁻¹. Monolayers were challenged in triplicate wells at a multiplicity of infection (MOI) of 100 : 1 (bacteria: epithelial cells). The plates were then incubated at 37°C and 5% CO₂ for 1 h. Following this, the monolayers were washed twice with SFM in order to remove nonadhered and loosely adhered bacteria. Cells were overlaid with SFM and further incubated for 30 min at 37°C and 5% CO₂. Monolayers containing associated (i.e. adhered and invaded) bacteria were lysed (in order to liberate the micro-organisms) with a 1 ml volume of 1% Triton-X-100 (Sigma) prepared in sterile PBS, for 5 min at room temperature. This 1 ml volume was serially diluted in PBS, and spread plates were prepared using selective agars.

Quantification of invasion

CaCo-2 cells were treated as previously described for the association assay. After nonadherent bacteria had been removed by washing with SFM, CaCo-2 cells were treated with gentamicin (Gibco) in order to quantify invasion. Gentamicin (an antibiotic) does not diffuse into CaCo-2 cells, so any externally adhered bacteria are rapidly killed but the viability of any invaded bacteria is not affected. Briefly, gentamicin was prepared at a concentration of 50 $\mu\text{g ml}^{-1}$ in SFM and added to wells, and monolayers were again incubated for 30 min at 37°C and 5% CO₂, before being washed twice with PBS to remove excess antibiotic. Epithelial cells were then lysed in order to liberate invaded pathogens and spread plates prepared as described earlier.

The quantity of associated/invaded pathogenic bacteria in SFM was assigned to 100%. Thus, percentage association/invasion was expressed relative to the control (in the absence of test material) as follows: (number of bacteria associated or invaded in the presence of test material/number of bacteria associated or invaded in the presence of DMEM alone) \times 100.

Mechanism of inhibition assays

In an attempt to determine whether test material interacted with either epithelial cells or bacteria or both, CaCo-2 cells and bacteria were separately preincubated with test material for 1 h at 37°C and 5% CO₂ prior to performing infection assays as described earlier. Three separate experiments were carried out for each of the micro-organisms ($n = 3$).

Statistical analysis

Experiments were carried out using three bacterial cultures ($n = 3$) for each treatment. Results were expressed as the mean \pm standard deviation (SD). Differences between inhibitory effects of each treatment were determined using the general linear models (GLM) function of SAS ver. 9.1.3 (SAS Institute, Cary, NC). Data were considered significantly different if $P < 0.05$.

Results

The compositional analysis of each test material was determined, along with the pH value of each whey product in its untreated (UT) and enzyme-modified (EM) form (Table 1). Test materials were dispersed in DMEM at a concentration of 40 mg ml⁻¹ prior to measuring pH, and the pH values of DMEM alone and egg albumin were 7.1 and 7.08, respectively. Enzyme-treatment was found to lower the pH of all whey products.

Table 1 Compositional analysis of whey products used in this study (%) and pH values of test materials before and after treatment with porcine pancreatic lipase

	SWPC80	AWPC80	SWPC35	WPI	WP	DW
Protein	75.5	78.2	34.3	86.6	12.5	13
Fat	8	7.7	3.4	0.1	1	1.8
Moisture	7.5	6.3	5.4	5.8	3.1	3.5
Ash	3	5.9	6.2	2.6	9.5	0.8
Lactose	6	1.9	50.7	4.9	73.9	80.9
pH UT	7.16	7.26	7.22	7.11	7.17	7.22
pH EM	6.48	6.56	6.58	6.56	7.01	6.97

SWPC80, sweet whey protein concentrate 80; AWPC80, Acid WPC80; SWPC35, sweet whey protein concentrate 35; WPI, whey protein isolate; WP, whey powder; DW, demineralized whey; UT, untreated; EM, enzyme-modified with porcine pancreatic lipase.

Test materials were not found to reduce viability of either CaCo-2 cells or bacteria at a concentration of 40 mg ml⁻¹ (data not shown).

Salm. Typhimurium

In the absence of test material, *Salm. Typhimurium* associated to and invaded CaCo-2 cells at levels of 10⁷ and 10⁶ CFU ml⁻¹ well⁻¹, respectively, and this is represented by the DMEM bars in Fig. 1a,b as 100% association/invasion. Of the untreated materials, all but AWPC80, DW and the protein control, egg albumin, significantly reduced association of *Salm. Typhimurium* to CaCo-2 cells ($P < 0.05$). UT SWPC80 was significantly more effective than the other materials, reducing association by c. 60% ($P < 0.05$). With one exception (WP), pretreatment of these whey products with PPL increased their ability to subsequently reduce *Salm. Typhimurium* association, although the reduction in association brought about by this treatment was not always significant (Fig. 1a).

In the invasion assays, of the untreated products examined, only SWPC80 and SWPC35 and to a lesser extent WPI reduced invasion significantly ($P < 0.05$). UT AWPC80 appeared to enhance invasion of *Salm. Typhimurium* into CaCo-2 cells (Fig. 1b). All EM whey products were significant inhibitors of invasion ($P < 0.05$), with EM-SWPC35 showing the greatest reduction (c. 75%). EM-SWPC80, EM-AWPC80, EM-WPI and EM-WP exhibited similar levels of activity, reducing invasion of this micro-organism into CaCo-2 cells by 40–50%. EM-DW was the least effective of the hydrolysates, but still reduced invasion by 26%.

E. coli O157:H7

Under the experimental conditions described, c. 10⁶ CFU ml⁻¹ well⁻¹ of *E. coli* O157:H7 cells associated





Figure 1 The Effect of DMEM (□), untreated whey products (□), enzyme-modified whey products (■) and egg albumin* on the (a) association and (b) invasion of *Salm.* Typhimurium to/into CaCo-2 cells. (Data = mean ± SD, $n = 3$). *Egg albumin is included in this figure for comparison only.




Figure 2 The effect of DMEM (□), untreated whey products (□), enzyme-modified whey products (■) and egg albumin* on the association of *E. coli* O157:H7 to CaCo-2 cells. (Data = mean ± SD, $n = 3$). *Egg albumin is included in this figure for comparison only.

to CaCo-2 monolayers in the absence of test material (representing 100% association, Fig. 2). The presence of all UT materials with the exception of DW significantly reduced association of *E. coli* O157:H7 with CaCo-2 cells ($P < 0.05$), as observed previously for association assays using *Salm.* Typhimurium. UT SWPC80 was again the most potent in this regard, reducing association by *c.* 60%. Interestingly, UT AWPC80 which did not reduce association of *Salm.* Typhimurium inhibited association of *E. coli* O157:H7 with CaCo-2 cells.

Pretreatment with PPL did not significantly increase the ability of the materials to reduce association ($P > 0.05$). An exception in this regard was EM-AWPC80, which showed the greatest reduction in association (>60% inhibition) of the materials studied. Also, it should be noted that egg albumin, the protein control,

was an effective inhibitor of association of *E. coli* O157:H7 with CaCo-2 cells, being only significantly less effective than EM-AWPC80 ($P < 0.05$).

Invasion of this micro-organism in the presence of the test materials was not examined here, as it has been reported that *E. coli* O157:H7 does not invade into all cell lines, but this does not necessarily mean that this bacterium is for tissue culture noninvasive (Oelschlaeger *et al.* 1994).

C. malonicus

In the absence of test material, this micro-organism showed similar levels of association with and invasion of CaCo-2 cells as those observed for *Salm.* Typhimurium (10^7 and 10^6 , respectively). Again these values represent




Figure 3 The effect of DMEM (■), untreated whey products (□), enzyme-modified whey products (■) and egg albumin* on the (a) association and (b) invasion of *C. malonaticus* to/into CaCo-2 cells (data = mean \pm SD, $n = 3$). *Egg albumin is included in this figure for comparison only.

100% association or invasion (Fig. 3a and Fig. 3b, respectively). UT AWPC80 increased association of *C. malonaticus* to CaCo-2 cells by *c.* 20% (Fig. 3a). No significant differences were noted between the potency of UT SWPC80, UT SWPC35, UT WPI and UT WP, but each of these whey products were found to be more effective than DMEM alone ($P < 0.05$), causing a 35–40% reduction in association. For the EM products, EM-SWPC80, EM-SWPC35 and EM-WPI were the most effective test materials, reducing association by 35–45%. EM-AWPC80, EM-WP and EM-DW had no significant effect on association of this bacterium to monolayers ($P > 0.05$). Egg albumin was found to significantly increase association of *C. malonaticus* by *c.* 50% to the CaCo-2 cell line and showed least inhibitory activity when compared to UT and EM whey products ($P < 0.05$).

For the invasion assays (Fig. 3b), UT AWPC80 appeared to increase invasion of *C. malonaticus* into epithelial cells by *c.* 55%. Of the UT whey products (Fig. 3b), the most effective inhibitors of invasion were UT WP and UT WPI, which showed greater reductions than UT SWPC80 and UT SWPC35 ($P < 0.05$). UT DW had no significant ($P > 0.05$) effect on invasion. However, all EM whey products greatly reduced invasion of *C. malonaticus* into CaCo-2 cells, causing a 2-log reduction when compared to DMEM alone ($10^6 \rightarrow 10^4$ CFU ml⁻¹ well⁻¹). EM-WP and EM-DW

were slightly less effective than other test materials, nevertheless causing a reduction of 92% and 88%, respectively (Fig. 3b). Egg albumin, the protein control, neither increased nor reduced invasion of this micro-organism into CaCo-2 cells.

Mechanism of action

EM-SWPC80 was the test material chosen for mechanistic experiments, as this whey product was found to be a very effective inhibitor of association of *Salm.* Typhimurium, *E. coli* O157:H7 and *C. malonaticus* to CaCo-2 cells. A reduction in association was observed for these micro-organisms when the bacteria and test material were added simultaneously to the test wells. However, with the exception of *C. malonaticus*, pretreatment of the micro-organisms with EM-SWPC80 generally did not reduce association, nor did pretreatment of the epithelial cells. The results of three separate experiments for each of the three micro-organisms are shown in Table 2.

Discussion

Adherence to the host's cell surface is a vital step in pathogenesis of gastrointestinal tract (GIT) infection as it prevents micro-organisms from being swept away by bulk fluid movement. Once a microbe adheres, it can readily

Table 2 CFU/ml/well of *Salm. Typhimurium*, *E. coli* O157:H7 and *C. malonaticus* associated to CaCo-2 Cells in the presence of EM-SWPC80 (40mg ml⁻¹) under varying assay conditions

Experiment	Maximum Association (i.e.100% association)*	EM-SWPC80 and bacteria added simultaneously	Bacteria pretreated with EM-SWPC80	CaCo-2 cells pretreated with EM-SWPC80
<i>Salm. Typhimurium</i>				
1	1.29 × 10 ⁷	4.3 × 10 ⁶ (33.3%)	1.53 × 10 ⁷ (0%)	1.77 × 10 ⁷ (0%)
2	1.43 × 10 ⁷	6.3 × 10 ⁶ (44.1%)	1.73 × 10 ⁷ (0%)	1.7 × 10 ⁷ (0%)
3	1.58 × 10 ⁷	4.85 × 10 ⁶ (61.8%)	2.03 × 10 ⁷ (0%)	1.78 × 10 ⁷ (0%)
<i>E. coli</i> O157:H7				
1	1.91 × 10 ⁶	6.2 × 10 ⁵ (32.3%)	1.78 × 10 ⁶ (93.2%)	1.69 × 10 ⁶ (88.5%)
2	1.6 × 10 ⁶	1.08 × 10 ⁶ (67.5%)	1.96 × 10 ⁶ (0%)	2.33 × 10 ⁶ (0%)
3	2.17 × 10 ⁶	1.6 × 10 ⁶ (73.4%)	1.56 × 10 ⁶ (0%)	2.5 × 10 ⁶ (0%)
<i>C. malonaticus</i>				
1	1.79 × 10 ⁷	7.8 × 10 ⁶ (43.6%)	7.1 × 10 ⁶ (39.2%)	1.82 × 10 ⁷ (0%)
2	9.8 × 10 ⁶	6.1 × 10 ⁶ (62.2%)	5.7 × 10 ⁶ (58.2%)	1.72 × 10 ⁷ (0%)

EM-SWPC80, Enzyme-modified sweet whey protein concentrate 80.

*Association of bacteria in DMEM alone is represented as 100%.

The reduction in association is shown in parentheses as a percentage and was calculated relative to the maximum association for that particular bacterial culture.

access nutrients, while being able to deliver toxins into host tissues before eventually penetrating these tissues (Acord *et al.* 2005). The aim of antiadhesion therapy is essential to reduce contact between pathogens and host tissues, by preventing or reversing adherence. Currently, some of the most efficient known inhibitors of adhesion are found in foodstuffs (Ofek *et al.* 2003). In the present study, the effect of native and EM whey products on association of *Salm. Typhimurium*, *E. coli* O157:H7 and *C. malonaticus* to human CaCo-2 cells was examined, along with their effects on the invasion of such cells by *Salm. Typhimurium* and *C. malonaticus*. Whey is currently described as a 'functional food', as it possesses a wide array of health benefits, and undenatured whey can provide high concentrations of intact proteins such as lactoferrin and immunoglobulins (Marshall 2004). The results described here indicate that, with the exception of AWPC80 and DW, most UT whey products were effective inhibitors of both association and invasion. However, the activity of these whey products was enhanced following hydrolysis with PPL, with EM-SWPC80 and EM-SWPC35 causing particularly high reductions.

Each whey product has varying levels of fat, protein, moisture, ash and lactose (Table 2). Sweet and acid WPCs have similar levels of protein and fat etc., yet the sweet WPCs were found to have greater anti-adhesion/invasion activity. WPI has almost no fat, yet still exhibited significant anti-adhesive and anti-invasion activity. WP and DW contain less protein and fat than the WPCs and WPI, but have a high content of lactose. In a report by Coppa *et al.* (2006), lactose from human milk (concentration not given) was not found to inhibit association of *E. coli* (serotype O119) or *Salm. Fyris* to CaCo-2 cells. In

the present study, inhibition of association and invasion of pathogenic bacteria by the whey products was generally more potent than that of egg albumin, suggesting the activity is not because of a nonspecific protein effect. Whey proteins are recognized as having superior biological activity to other proteins, and the activity of peptides encrypted within such proteins usually remains latent until they are subjected to proteolytic action of enzymes (Sinha *et al.* 2007). The crude PPL used in the present study is known to contain both proteases and lipases, and it may be that enzyme pretreatment of the whey products listed here releases species such as specific peptides or free fatty acids that are latent within the untreated material. The observation that potency of these materials was not diminished after enzyme treatment may indicate that the active components could survive passage through the gut. This would be a favourable characteristic for using whey products to prevent/treat infection of the GIT.

One potential hypothesis as to why high levels of inhibition were observed for sweet WPCs may be because of the presence of glycomacropeptide (GMP). GMP, also referred to as the 'casein macropeptide' is present in sweet (rennet) whey as a result of the action of the chymosin enzyme on κ -casein during the cheese-making process (Marshall 2004). This peptide is known to inhibit bacterial and viral adhesion (Kawasaki *et al.* 1993; Simon 1996) and is capable of binding to *E. coli* O157:H7 and *Salmonella* Enteridis when in its sialylated form (Nakajima *et al.* 2005). In a study by Brück *et al.* (2006), the effect of GMP on the association of enteropathogenic *E. coli* (EPEC) and *Salm. Typhimurium* to CaCo-2 cells was examined, and they found that undigested GMP (0.25 mg ml⁻¹) reduced association of these bacteria.

They also reported that enzyme digestion with pepsin and pancreatin increased the antiadhesive activity of GMP. It was concluded that fragments produced by digestion with both pancreatin and pepsin digestion of GMP were most potent for *Salm. Typhimurium*, but peptides liberated from digestion with pepsin alone significantly reduced association levels of EPEC ($P < 0.05$ – 0.001). However, isolated and purified GMP is an expensive resource. Our study has shown that native sweet WPC products, which are inexpensive and readily available, are effective inhibitors of bacterial association and invasion *in vitro*. Enzyme-modifying such material with PPL is also a relatively inexpensive and straightforward process, and enhances the anti-adhesive and anti-invasive activity of sweet WPCs.

In a study by de Araujo and Giugliano (2000), it was reported that whey produced from human milk (0.97 mg ml^{-1}) inhibited association of diffusely adherent *E. coli* (DAEC) and enteroaggregative *E. coli* (EAEC) to HeLa cells by 9% and 16%, respectively. In a subsequent study by the same researcher in 2001, the free secretory component ($105 \mu\text{g ml}^{-1}$) and lactoferrin ($157 \mu\text{g ml}^{-1}$) isolated from human milk, inhibited adherence of EPEC to HeLa cells by 32% and 4.5%, respectively. The immunoglobulin fraction was also found to reduce adherence of these bacteria to HeLa cells (de Araujo and Giugliano 2001). da Motta Willer *et al.* (2004) found that whey from human milk (2.8 mg ml^{-1}) could reduce adhesion of *Shigella* strains to HeLa cells by at least 40%, and invasion was reduced by more than 50%. It was also reported that human lactoferrin (0.3 mg ml^{-1}) reduced invasion by 50–65% (da Motta Willer *et al.* 2004). It is possible that the equivalent components of bovine milk/whey could inhibit association and/or invasion of pathogens to epithelial cells.

Nonprotein components of whey include oligosaccharides and lipids, including sphingolipids (Shoaf *et al.* 2006). Certain oligosaccharides are similar in structure to receptor sites (recognized and adhered to by pathogens) coating epithelial cells of the intestine. Thus, oligosaccharides possibly act as molecular receptor decoys, competitively inhibiting microbial adherence. Conversely, instead of pathogens adhering to the host cell surfaces, they would bind to the decoy oligosaccharides and be displaced from the GIT (Shoaf *et al.* 2006). Another potential hypothesis as to why the whey products used in this study inhibited association of pathogenic bacteria to epithelial cells is the presence of oligosaccharides. The oligosaccharide content of the whey products used in our study may be contributing to the anti-adhesive activities of these materials. Another characteristic of the test materials used in this study, which should be taken into consideration, are the pH

values of individual products when in solution. Treatment with PPL reduced the pH of all test materials, because of the liberation of free fatty acids and amino acids during hydrolysis. It is possible that the fimbrial protein structures of pathogenic micro-organisms are influenced by different pH values (close to their pI values), reducing the capability of bacteria to adhere to epithelial cells (Lehto and Salminen 1997). This may be a contributing factor in the reduction in association of *Salm. Typhimurium*, *E. coli* O157:H7 and *C. malonicus* to CaCo-2 cells in the presence of whey products, which was observed in the results reported here. Preliminary results have shown that inhibition of association is greatest when EM-SWPC80 and bacteria are added simultaneously to monolayers (Table 2). However, whether this inhibitory effect is caused by proteins, fats, pH, oligosaccharides, immunoglobulins, lactobacilli, etc. present in whey or a combination of these is not yet clear, and the exact mechanism of action of these products to prevent association/invasion of pathogenic bacteria to epithelial cells remains to be elucidated, but it is not unreasonable to speculate that the inhibitory effect is because of the presence of GMP.

Although CaCo-2 cells are regarded as one of the best *in vitro* models of mature enterocytes, such cell systems have their limitations (Giannasca *et al.* 1996), such as the absence of host factors (e.g. mucus barriers, immune z-factors). In addition, cell culture models do not possess other host cells that would normally be present *in vivo* e.g. inflammatory cells (Finlay and Falkow 1997). Nevertheless, the results of the present study are positive from a cell culture perspective.

Individuals most at risk from foodborne infection include newborns, elderly people and those who are immunocompromised (Sprong *et al.* 1999). Treatments for infection caused by foodborne pathogens such as those caused by *E. coli* O157:H7 is for the most part supportive, as practitioners are reluctant to prescribe antibiotics because of the risk of complications, such as acute renal failure in patients with HUS (Meng and Doyle 1998). Also, the emergence of multidrug-resistant strains of *Salm. Typhimurium* has led to a limited number of treatment options, in particular for invasive infections (Meng and Doyle 1998). Thus, the use of antibiotics to treat foodborne illness is no longer desirable because of the complications which are likely to occur. These include incidences of drug-resistant strains and the potential for chronic toxicity (Lin *et al.* 2007). As a result, alternative approaches to preventing infections of the GIT have been sought. Whey products, in either their UT or EM form may be suitable alternatives to treat or preferably prevent illness caused by foodborne pathogens.

Conclusion

Both UT and EM whey products are effective inhibitors of association of *Salm. Typhimurium*, *E. coli* O157:H7 and *C. malonaticus* to CaCo-2 cells and may be suitable for *in vivo* use to prevent and/or treat GIT infection caused by foodborne pathogens.

Acknowledgements

This work was supported by the National Development Plan (NDP), with a grant from the Food Institution Research Measure (FIRM). We thank Carbery Ireland for supplying the whey products used throughout this study, and also we thank Maeve O'Connor for carrying out compositional analysis on these whey products.

References

- Acord, J., Maskell, J. and Sefton, A. (2005) A rapid microplate method for quantifying inhibition of bacterial adhesion to eukaryotic cells. *J Microbiol Methods* **60**, 55–62.
- de Araujo, A.N. and Giugliano, L.G. (2000) Human milk fractions inhibit the adherence of diffusely adherent *Escherichia coli* (DAEC) and enteroaggregative *E. coli* (EAEC) to HeLa cells. *FEMS Microbiol Lett* **184**, 91–94.
- de Araujo, A.N. and Giugliano, L.G. (2001) Lactoferrin and free secretory component of human milk inhibit the adhesion of enteropathogenic *Escherichia coli* to HeLa cells. *BMC Microbiology* **1**, 25–30.
- Bezirtzoglou, E., Maipa, V., Voidarou, C., Tsiotsias, A. and Papapetropoulou, M. (2000) Food-borne intestinal bacterial pathogens. *Microb Ecol in Health Dis* **2**(suppl.), 96–104.
- Brück, W.M., Kelleher, S.L., Gibson, G.R., Graverholt, G. and Lonnerdal, B.L. (2006) The effects of α -lactalbumin and glycomacropeptide on the association of CaCo-2 cells by enteropathogenic *Escherichia coli*, *Salmonella typhimurium* and *Shigella flexneri*. *FEMS Microbiol Lett* **259**, 158–162.
- Coppa, G.V., Zampini, L., Galeazzi, T., Facinelli, B., Ferrante, L., Capretti, R. and Orazio, G. (2006) Human milk oligosaccharides inhibit the adhesion to CaCo-2 cells of diarrheal pathogens: *Escherichia coli*, *Vibrio cholerae* and *Salmonella typhi*. *Paediatr Res* **59**, 377–382.
- Finlay, B.B. and Falkow, S. (1997) Common themes in microbial pathogenicity revisited. *Microbiol Mol Biol Rev* **61**, 136–169.
- Fogh, J., Fogh, J.M. and Orfeo, T. (1977) One hundred and twenty-seven cultured human tumour cell lines producing tumors in nude mice. *J Natl Cancer Inst* **59**, 221–226.
- Giannasca, K.T., Giannasca, P.J. and Neutra, M.R. (1996) Adherence of *Salmonella typhimurium* to CaCo-2 cells: identification of a glycoconjugate receptor. *Infect Immun* **64**, 135–145.
- Gu, L., Wang, H., Guo, Y.-L. and Zen, K. (2008) Heparin blocks the adhesion of *E. coli* O157:H7 to human colonic epithelial cells. *Biochem Biophys Res Commun* **369**, 1061–1064.
- Gurtler, J.B., Kornacki, J.L. and Beuchat, L.R. (2005) *Enterobacter sakazakii*: a coliform of increased concern to infant health. *Int J Food Microbiol* **104**, 1–34.
- Halpin, R.M., O'Connor, M.M., McMahon, A., Boughton, C., O'Riordan, E.D., O'Sullivan, M. and Brady, D.B. (2008) Inhibition of adhesion of *Streptococcus mutans* to hydroxylapatite by commercial dairy powders and individual milk proteins. *Eur Food Res Technol* **227**, 1499–1506.
- Hendricks, H., van Asten, A., Koninx, J., Kok, W., van der Zeijst, B. and van Dijk, J. (1996) Interactions between *Salmonella* Enteridis and the enterocyte-like human carcinoma cell line CaCo-2: effects of nutrients on the nutritional value of legume diets. In *Luxembourg: Office Official Publications European Communities* Bardocz, S., Nekrep, F.V. and Pustazi, A. ed. pp. 137–139. Brussels: European Commission.
- International Dairy Federation (1987) *Standard 9C: Determination of Fat Content of Dried Milk, Dried Whey, Dried Buttermilk and Dried Butter*. Brussels: International Dairy Federation.
- International Dairy Federation (1993a) *Milk, Determination of the Nitrogen Content: II. Block Digestion Method (standard 20B)*. Brussels: International Dairy Federation.
- International Dairy Federation (1993b) *Dried Milk and Cream-Determination of Water Content*. Brussels: International Dairy Federation.
- Iversen, C., Lehner, A., Mullane, N., Marugg, J., Fanning, S., Stephan, R. and Joosten, H. (2007) Identification of “Cronobacter” spp. (*Enterobacter sakazakii*). *J Clin Microbiol* **45**, 3814–3816.
- Kandhai, M.C., Reij, M.W., Gorris, L.G.M., Guillaume-Gentil, O. and van Schothorst, M. (2004) Occurrence of *Enterobacter sakazakii* in food production environments and households. *Lancet* **363**, 39–40.
- Kaper, J.B., Nataro, J.P. and Mobley, H.L.T. (2004) Pathogenic *Escherichia coli*. *Nat Rev* **2**, 123–140.
- Kawasaki, Y., Isoda, K., Shinmoto, H., Tanimoto, M., Dosako, S., Idota, T. and Nakajima, I. (1993) Inhibition by κ -casein glycomacropeptide and lactoferrin of influenza virus hemagglutination. *Biosci Biotechnol Biochem* **57**, 1214–1215.
- Kim, K.-P. and Loessner, M.J. (2008) *Enterobacter sakazakii* invasion in human intestinal CaCo-2 cells requires the host cell cytoskeleton and is enhanced by disruption of tight junctions. *Infect Immun* **76**, 562–570.
- Koninx, J.F.J.G. (1995) Enterocyte-like CaCo-2 cells as a tool to study lectin interaction. In *Lectins: Biomedical Perspectives* ed. Pusztai, A. and Bardocz, S. pp. 81–10. London: Taylor and Francis.
- Lehto, E.M. and Salminen, S.J. (1997) Inhibition of *Salmonella typhimurium* adhesion to CaCo-2 cell cultures by *Lactobacillus* strain GG spent culture supernatant: only a pH effect? *FEMS Immunol Med Microbiol* **18**, 125–132.

- Lin, W.-H., Yu, B., Lin, C.-K., Hwang, W.-Z. and Tsen, H.-Y. (2007) Immune effect of heat-killed multistrain of *Lactobacillus acidophilus* against *Salmonella typhimurium* invasion to mice. *J Appl Microbiol* **102**, 22–31.
- Malkomesius, P.E. and Nehring, K. (1951) Chemische Untersuchung von Futtermitteln. In *Handbuch der landwirtschaftlichen Versuchs- und Untersuchungsmethodik, band 3* ed. Herrmann, R. pp. 15, 25. Berlin, Germany: Naumann Verlag.
- Mange, J.-P., Stephan, R., Borel, N., Wild, P., Kim, K.S., Pospischil, A. and Lehner, A. (2006) Adhesive properties of *Enterobacter sakazakii* to human epithelial and brain microvascular endothelial cells. *BMC Microbiol* **6**, 58–68.
- Marshall, K. (2004) Therapeutic applications of whey protein. *Altern Med Rev* **9**, 136–156.
- Meisel, H. (1998) Overview on milk protein-derived peptides. *Int Dairy J* **8**, 363–373.
- Meng, J. and Doyle, M.P. (1998) Emerging and evolving microbial foodborne pathogens. *Bull Inst Pasteur* **96**, 151–164.
- da Motta Willer, E., de Lourenco Lima, R. and Giugliano, L.G. (2004) *In vitro* adhesion and invasion inhibition of *Shigella dysenteriae*, *Shigella flexneri* and *Shigella sonnei* clinical strains by human milk proteins. *BMC Microbiol* **4**, 18–25.
- Mullane, N.R., Iversen, C., Healy, B., Walsh, C., Whyte, P., Wall, P.G., Quinn, T. and Fanning, S. (2007) *Enterobacter sakazakii*: an emerging bacterial pathogen with implications for infant health. *Minerva Paediatr* **59**, 137–148.
- Nakajima, K., Tamura, N., Kobayashi-Hattori, K., Yoshida, T., Hara-Kudo, Y., Ikedo, M., Sugita-Konishi, Y. and Hattori, M. (2005) Prevention of intestinal infection by glycomacropeptide. *Biosci Biotechnol Biochem* **69**, 2294–2301.
- Oelschlaeger, T.A., Barrett, T.J. and Kopecko, D.J. (1994) Some structures and processes of human epithelial cells involved in uptake of enterohemorrhagic *Escherichia coli* O157:H7 strains. *Infect Immun* **62**, 5142–5150.
- Ofek, I., Hasty, D.L. and Sharon, N. (2003) Anti-adhesion therapy of bacterial diseases: prospects and problems. *FEMS Immunol Med Microbiol* **38**, 181–191.
- Pinto, M., Robine-Leon, S., Appay, M.-D., Kedinger, M., Triadou, N., Dussaulx, E., Lacroix, B., Simon-Assmann, P. et al. (1983) Enterocyte-like differentiation and polarisation of the human colon carcinoma cell line CaCo-2 in culture. *Biol Cell* **47**, 323–330.
- Rodrigue, D.C., Tauxe, R.V. and Rowe, B. (1990) International increase in *Salmonella enteridis*: a new pandemic? *Epidemiol Infect* **105**, 21–27.
- Shoaf, K., Mulvet, G.L., Armstrong, G.D. and Hutkins, R.W. (2006) Prebiotic galactooligosaccharides reduce adherence of enteropathogenic *Escherichia coli* to tissue culture cells. *Infect Immun* **74**, 6920–6928.
- Simon, P.M. (1996) Pharmaceutical oligosaccharides. *Drug Discov Today* **1**, 522–528.
- Sinha, R., Radha, C., Prakash, J. and Kaul, P. (2007) Whey protein hydrolysate: functional properties, nutritional quality and utilisation in beverage formulation. *Food Chem* **101**, 1484–1491.
- Sprong, R.C., Hulstein, M.F. and Van der Meer, R. (1999) High intake of milk fats inhibits intestinal colonisation of *Listeria* but not *Salmonella* in rats. *J Nutr* **129**, 1382–1389.
- Sugita-Konishi, Y., Sakanaka, S., Sasaki, K., Juneja, L.R., Noda, T. and Amano, F. (2002) Inhibition of bacterial adhesion and *Salmonella* infection in BALB/c mice by sialyloligosaccharides and their derivatives from chicken egg yolk. *J Agric Food Chem* **50**, 3607–3613.
- Tang, P., Foubister, V., Pucciarelli, G. and Finlay, B.B. (1993) Methods to study bacterial invasion. *J Microbiol Methods* **18**, 227–240.
- WHO (2005) *Global Salm-Surv. Top 15 Salmonella Serotype List from Each Country*. http://thor.dfvf.dk/pls/portal/GSS.COUNTRY_DATA_SET_REP.show.